

# A victory for all mankind

Brought about by hundreds of thousands of health workers from all over the world, eradication is a triumph of international cooperation and of preventive medicine

by D. A. Henderson



For centuries, variola virus stalked the world with impunity causing unmeasured suffering, death and blindness. Today it is confined to glass vials kept under high security in six laboratories. To those who vividly recall the suffering of its victims, to health staff who so diligently conducted vaccination programmes, to quarantine inspectors who carefully examined vaccination certificates, it is difficult to believe that "sporadic" cases are not still occurring somewhere. However, convincing data, diligently assembled by tens of thousands of health staff, now demonstrate that smallpox is a disease which can be consigned to history—the first disease ever eradicated by man.

It was little more than two decades ago, in 1958, that the Soviet Union proposed to the Eleventh World Health Assembly that the countries of the world cooperate in a globally coordinated effort to eradicate smallpox. In that year, 63 countries reported 280,000 cases. This is itself a substantial number of cases but, because of incomplete reporting, it would need to be multiplied by 100 or more to describe the true magnitude of the problem. Data regarding deaths are likewise incomplete, but it is reasonable to assume that at least 20 per cent of those infected died and tens of thousands were permanently blinded. There was, and is, no treatment for smallpox; prevention through vaccination was the only way to cope with this disease.

After further study of the Soviet proposal, the Assembly in 1959 agreed that a global smallpox eradication programme should be undertaken as a matter of urgency. It was thought that, by "vaccinating or revaccinating 80 per cent of the

population within a period of four to five years", smallpox could be eradicated from endemic areas.

During the following eight years, WHO encouraged governments to undertake programmes, solicited contributions of vaccine, encouraged and coordinated studies of vaccine strains, and assisted laboratories in starting vaccine production. Many countries started programmes and some succeeded in interrupting transmission. But not all did or could embark on eradication with their own limited resources. Contributions were far less than the funds required and many countries which interrupted transmission were reinfected by their neighbours. Of greater concern was the discovery that, even when vaccinations were administered to 80 per cent of a population, smallpox often persisted. Health authorities became discouraged and pessimistic. It was one thing to eliminate the disease from countries with a developed health infrastructure, but could this be done in the numerous countries, so recently independent, whose limited health services scarcely extended beyond their urban centres? And could these many countries coordinate their efforts sufficiently for smallpox to be eliminated from large geographical areas?

It was a concerned and less confident World Health Assembly which in 1966 decided that WHO should undertake an intensified eradication programme. Approximately US \$2.5 million was included in the Organization's regular budget to provide for overall programme coordination and for assistance to those countries requiring it. This was a small sum indeed to provide support in some 50 countries with a population of more than one thousand million persons. But it represented almost five per cent of WHO's total budget that year. Hopes were expressed that more substantial vol-

*This was smallpox. The world will never again witness the suffering that was caused by the*

Smallpox is dead  
Cover design  
by Peter Davies

World Health  
May 1980: 3-5

IX ISSN 0043-8502

World Health is the official  
magazine of the World Health O

## Editor:

John Bland

## Deputy Editor:

Christiane Viedma

## Art Editor:

Peter Davies

## News Page Editor:

Lalit Thapalyal

World Health appears in Arabic, French, German, Italian, Persian, Russian and Spanish.

Articles and photographs not reproduced provided credit to the World Health Organization; articles do not necessarily reflect views.

World Health, WHO,  
Av. Appia, 1211 Geneva 27, Switzerland

## Contents

### A victory for all mankind

by D. A. Henderson

### A windfall for development

by James Magee

### A job well done

by Marcella Davies

### An effort of will

by V.T. Herat Gunaratne

### India's "war plan"

by Jitendra Tuli

### A goddess defied

by Arun M. Chacko

### How much did it cost?

### Towards the year 2000

by Halfdan Mahler

### Ramses V: earliest known victim

by Donald R. Hopkins

### Decline and Fall

of the Smallpox Empire

### Can we stop smallpox vaccination?

by Isao Arita

### The closing stages

by Svetlana Marennikova

### How can we be sure?



that all countries would give the programme a high priority. Some delegates optimistically proposed setting a 10 years goal for eradication, but most believed realistically that better control of smallpox was the best that could be expected.

The intensified programme began in January 1967. That year, 46 countries recorded 131,697 cases, a number representing perhaps one per cent of the true number of cases. Four endemic areas were present. A major reservoir was Africa where virtually all countries south of the Sahara were infected. A second important reservoir of smallpox was in Asia, extending from Bangladesh through India, Nepal, Pakistan and Afghanistan. The third was the Indonesian archipelago and the fourth was Brazil, itself comprising half a continent.

Of primary concern as the programme began was the need for sufficient vaccination devices and vaccine—the guns and bullets of the campaign. Large contributions were made initially by the Soviet Union and the USA; eventually 26 countries became contributors. A detailed manual on vaccine production was produced and vaccine batches were routinely tested to ensure that they met international standards. By 1971, all vaccine in use in the programme met accepted standards and by 1973, fully 80 per cent was being produced in the endemic countries, some of which supplied vaccine to others. In 1967, the jet injector was introduced in programmes throughout the countries of western and central Africa and Brazil. In 1968, field studies conducted by WHO showed that the newly developed bifurcated needle could be used to administer vaccine by a new technique—multiple puncture vaccination. The

bifurcated needle soon became the standard method for vaccination.

A second concern from the start was the question of an appropriate strategy. Mass vaccination designed to reach 80 or even 100 per cent of a population had succeeded only in some smaller countries and those with more developed health services. For most of the still endemic countries, a different strategy was required. The decision to emphasize surveillance as an important component of the strategy proved to be the much-needed critical breakthrough. The new strategy called for a systematic two- to three-year vaccination campaign designed to reach 80 per cent of the population in each country. During this time, it was planned for a nationwide reporting system to be developed which would be sensitive enough to detect such smallpox foci as remained and to eliminate them.

Soon after the programme began, it was discovered first in Nigeria, then in Indonesia and Brazil, that effective reporting systems could be developed in months rather than years. By isolating the patients and vaccinating their contacts, outbreaks could be rapidly contained. Even in areas where vaccination coverage was poor, smallpox transmission could often be stopped quickly. So now increasing emphasis was placed on the surveillance-containment component of the strategy. Special surveillance teams were recruited and trained. They visited each health unit in an area to ensure that each week it submitted a report indicating the number of cases seen. When cases were reported, the teams worked with local health staff to contain the outbreaks, and they visited schools and public places to inquire about

*The end of smallpox in Asia meant that more resources could be made available in Ethiopia. The net was closing on the last places on earth where the disease lurked. (WHO) E. Shafa*

rumours of smallpox. A special "WHO Recognition Card" showing a picture of a smallpox patient was printed and distributed to help them in their search.

The first campaigns started in 1967 and, by 1969, all countries except Ethiopia had started eradication programmes. Ethiopia's programme began in 1971. In 20 countries of western and central Africa, a USA-assisted programme of smallpox eradication and measles control succeeded in eliminating smallpox in just three and a half years. Brazil's last case was detected in 1971 and Indonesia's in 1972. By the summer of 1973, smallpox transmission had been interrupted throughout the whole of Africa, except in Ethiopia. That country plus five countries in Asia remained as the only smallpox infected countries.

However, India, Pakistan and Bangladesh, with a population of more than 700 million persons, presented a special problem. Surveillance-containment measures such as had been successful in Africa and South America proved far less effective in these densely populated areas where people travel frequently and far. A different approach was required. During the summer of 1973, Indian health authorities with WHO staff planned a different surveillance strategy. All health personnel were asked to undertake an intensive search of each village, and later each house, to detect smallpox cases. This would need to be done rapidly to be successful and so the search was planned to be completed during a seven- to ten-

day period. In heavily infected areas this was done each month and, in less infected areas, every second or third month. During the first search in October 1973, thousands of unreported cases were detected. But once discovered, the outbreaks could be contained.

Through careful planning, training and assessment, the searches became increasingly thorough. Between searches, surveillance teams visited schools and markets seeking to learn about cases of smallpox. Containment measures were strengthened. As the numbers began to decrease, a reward for the detection of a case was offered to the first person reporting it and to the health worker investigating it. Similar approaches were soon employed in Bangladesh, Pakistan and Nepal. On 16 October 1975, only two years after the start of this new strategy, the last case of smallpox occurred in Asia—on Bhola Island in Bangladesh. This marked the end of variola major, the severe form of the disease. As 1975 drew to a close, only Ethiopia remained with smallpox. There, variola minor was prevalent, a form of smallpox which killed only one per cent of its victims in contrast to variola major which killed 20 per cent or more.

Ethiopia's programme, besides being the last to begin, faced incredibly difficult problems. Although the country is larger than France and Spain together, the available staff numbered little more than 100 persons. There was almost no health infrastructure. Roads were few and it was estimated that half the population lived more than a day's walk from any accessible road. Civil war and famine compounded the problems. However, with the interruption of smallpox transmission in Asia, more resources could be made available to Ethiopia. Village residents were recruited and trained to serve as surveillance workers and vaccinators; more transport was provided; helicopters were laid on to facilitate supervision. Less than one year later, on August 1976, the last case occurred here.

Unfortunately, coincident with the last outbreak in Ethiopia, smallpox was introduced into neighbouring Somalia. Before fully effective surveillance systems could be established, nomads disseminated the disease throughout the southern part of the country. In May 1977, a national emergency was declared, additional staff were recruited and special assistance was provided by

ment and vaccination programme continued throughout the summer months. More than 3,000 cases occurred, the last of them on 26 October 1977.

Nevertheless, surveillance teams and search workers continued for two years and more in their efforts to discover cases. In many countries, a reward was

## A windfall for development

Already the international community is starting to collect part of the health legacy—and it is likely to be huge—that smallpox eradication will bequeath. Calculations indicate that in the post-smallpox era a sum of nearly US \$1,000 million annually will be released, or some \$10,000 million over a decade.

The cost to the world of a disease like smallpox included production or purchase of vaccine, maintenance of vaccination programmes, the treatment of vaccination complications, spending to maintain national surveillance and frontier controls, and the cost involved in handling the emergencies caused by sudden outbreaks. Thus in the United Kingdom, an outbreak sparked by an imported case in 1961 involved a bill for an estimated \$3.6 million. United States experts calculate that smallpox protection was costing the American taxpayer about \$150 million a year—or about half the total cost of the global eradication programme which was just over \$300 million.

The release of money that has hitherto been tied up for smallpox could have massive impact—provided it is diverted to development programmes. In one recent year, 1975, total funds available for the health sector (including water supplies) as bilateral aid from the Organization for Economic Cooperation and Development (OECD) and from the European Economic Community (EEC), or as multilateral aid via the UN specialized agencies, amounted to \$1,500 million. The new resources could therefore make a massive addition to the funds available.

In the view of WHO, these funds would have their greatest strategic impact in the area of primary health care. They could be applied to a combination of health development activities, including clean water and sanitation, immunization programmes, action to promote correct and adequate nutrition, particularly for infants and young children, maternal and child care, and programmes for the supply of oral rehydration kits to combat diarrhoeal diseases and to ensure the availability of essential drugs to all who need them.

offered and tens of thousands of cases of chickenpox, measles and other rash diseases were reported by villagers in hopes of collecting a reward. Thousands of specimens were taken and sent to WHO Diagnostic Centres in Moscow and Atlanta, USA. None proved to be smallpox. For the disease to persist in a population, it must be passed from one person to another in a continuing chain of infection. Since there is no animal reservoir and no asymptomatic carrier (person having the disease but showing no symptoms), each person who is a link in the chain must experience the characteristic illness. During the programme, it was found that smallpox never persisted for more than eight months in a country without being detected by the surveillance network. Continuing search during a two-year period, that is, three times this eight-month period, provided additional assurance that the spread of smallpox had stopped.

To confirm that transmission had been interrupted, specially constituted WHO International Commissions visited each country. In 1978, the Director-General of WHO appointed a Global Commission to review the experience in all countries. The Commission completed its work on 9 December 1979, at which time it agreed that there was sufficient documentation to certify worldwide eradication.

The eradication of smallpox means that vaccination everywhere can be stopped and that travellers will no longer need vaccination certificates. The savings throughout the world are estimated to be \$1,000 million annually. The cost of this achievement was only \$112 million in international assistance, or \$9 million spent annually during the 13-year period 1967-1979. This includes all funds spent by WHO plus bilateral and multilateral contributions from 42 countries. It is estimated that globally the endemic countries spent perhaps twice this amount but, in fact, few spent much more than they had hitherto been spending on never-ending smallpox control programmes.

Now the chapter entitled "smallpox" is closed—let us hope for ever. The achievement of eradication is a victory for hundreds of thousands of health workers in many different countries. WHO staff alone were drawn from 73 different countries. It is a triumph of international cooperation and of preventive medicine. With coordination, cooperation, dedication and imagination it is clear that much more can be achieved